



## **Bioelectricity production and microbial communities in microbial fuel cell powered by macroalgal biomass**

**Zhao, Nannan; Jiang, Yinan ; Alvarado-Morales, Merlin; Treu, Laura; Angelidaki, Irini; Zhang, Yifeng**

*Published in:*  
Book of Abstracts, Sustain 2017

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Zhao, N., Jiang, Y., Alvarado-Morales, M., Treu, L., Angelidaki, I., & Zhang, Y. (2017). Bioelectricity production and microbial communities in microbial fuel cell powered by macroalgal biomass. In *Book of Abstracts, Sustain 2017* [R-2] Technical University of Denmark.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Bioelectricity production and microbial communities in microbial fuel cell powered by macroalgal biomass

Nannan Zhao<sup>1</sup>, Yinan Jiang<sup>2</sup>, Merlin Alvarado-Morales<sup>2</sup>, Laura Treu<sup>2</sup>, Irini Angelidaki<sup>2</sup>, Yifeng Zhang<sup>\*3</sup>

1: Department of Environmental Engineering, Technical University of Denmark, DK-2800 Lyngby, Denmark

\*Email address: [yifz@env.dtu.dk](mailto:yifz@env.dtu.dk)



The potential of macroalgae *Laminaria digitata* as substrate for bioelectricity production was examined in a microbial fuel cell (MFC). A stable voltage of 0.6 V (at 1000 ohm) was achieved without any lag time due to the high concentration of glucose and mannitol in the hydrolysate, and it lasted for over 35 days. Total TCOD removal efficiency remained high and reached over 95% at the end of the test. However, the coulombic efficiency was low approx. 12%. The volatile fatty acids analysis implied that glucose and mannitol were degraded through isobutyrate as intermediate. During the operation, pH in anode and cathode exhibited reverse variation because of the proton accumulation in anode. The 16S rRNA gene high throughput sequencing analysis of anodic biofilm revealed complex microbial composition dominated by *Bacteroidetes* (39.4%), *Firmicutes* (20.1%), *Proteobacteria* (11.5%), *Euryarchaeota* (3.1%), *Deferribacteres* (1.3%), *Spirochaetes* (1.0%), *Chloroflexi* (0.7%), *Actinobacteria* (0.5%), and others (22.4%). The predominance of *Bacteroidetes*, *Firmicutes* and *Proteobacteria* demonstrated their importance for substrate degradation and simultaneous power generation. These results demonstrate that macroalgae hydrolysate can be used as a renewable carbon source of microbial electrochemical systems for various environmental applications.